₹¶	<b>Routine Vaccine Administration Guide</b> I Please refer to the DHW Publicly Funded Vaccine Schedule here			Publicly Funded Vaccines in Nova Scotia Childhood, Starting in Infancy (By Age)				
	VACCINE	PRODUCT NAME	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SI	ZE
	DTaP-IPV-Hib 🌗	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g	2 separate needles at
2 MONTHS	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	1-inch	heedles at least 1-inch apart
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable	
4. months	DTaP-IPV-Hib 🌗	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g	2 separate needles at least 1-inch apart
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	1-inch	
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable	
6	DTaP-IPV-Hib 🌗	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch	
MONTHS	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable	
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 -	2 separate needles at least 1-inch apart
<b>12</b> MONTHS	Meningococcal group C conjugate	NeisVac-C Menjugate	Vastus lateralis (thigh)	Intramuscular	0.5 mL	NeisVac-C (No) Menjugate (Yes or pre-filled)	25 g 1-inch	
	MMRV (measles, mumps, rubella, varicella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine- specific diluent)	25 g 5/8-inc	h
10	DTaP-IPV-Hib 🌗	Pediacel	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch	
<b>18</b> молтнs	MMRV (measles, mumps, rubella, varicella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine- specific diluent)	25 g 5/8-inc	h
<b>4-6</b> years	Tdap-IPV 🌗	Adacel Polio Boostrix Polio	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch	
ANNUALLY, FROM 6 MONTHS OF AGE ONWARD	Influenza 1st-time recipients aged 6 months – 9 years: 2 doses (0, 4 weeks)	Fluzone Flulaval Tetra Afluria (>5 years old only)	6-12 months old: vastus lateralis (thigh) Older than 12 months: deltoid	Intramuscular	0.5 mL	No	25 g 1-inch	

U vaccines with similar names that may lead to vaccine errors. i Products are subject to change based on availability.

#### **Catch-up (Resuming or Starting Immunizations)**

- Disruptions in vaccine schedules do not require that a series be re-started, regardless of the length of time between doses.
- For catch-up schedules, please refer to the Canadian Immunization Guide for age-specific recommendations:

# Additional Publicly Funded Vaccines for Patients at High-Risk of Vaccine-preventable Disease

Patients at high-risk of vaccine-preventable diseases are eligible for other vaccines based on **condition-specific guidance.** 



- Less than 7 years old; 7 - 17 years old; 18 years or older



	PRODUCT NAME	TOTAL DOSES	AGES	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZE
DTaP-IPV-Hib ()	Pediacel	4	2, 4, 6, 18 months	< 12 months: Vastus lateralis (thigh) Older than 12 months: Deltoid	Intramuscular	0.5 mL	No	25 g 1 inch
ROTAVIRUS	RotaTeq	3	2, 4, 6 months	Orally administered	Oral	2 mL	No	Not applicable
PNEUMOCOCCAL CONJUGATE (PNEU-C-13)	Prevnar 13	3	2, 4, 12 months	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1 inch
MENINGOCOCCAL GROUP C CONJUGATE	NeisVac-C Menjugate	1	12 months	Vastus lateralis (thigh)	Intramuscular	0.5 mL	NeisVac-C (No) Menjugate (Yes)	25 g 1 inch
MMRV (MEASLES, MUMPS, RUBELLA, VARICELLA)	Priorix Tetra ProQuad	2	12, 18 months	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine- specific diluent)	25 g 5/8 inch
Tdap-IPV	Adacel Polio Boostrix Polio	1	4 -6 years	Deltoid	Intramuscular	0.5 mL	No	25 g 1 inch
INFLUENZA	Fluzone Flulaval Tetra Afluria (>5 years old only)	Annually; 1st-time recipients aged 6 months – 9 years: 2 doses (0, 4 weeks)	From 6 months of age onward	6-12 months old: vastus lateralis (thigh) Older than 12 months: deltoid	Intramuscular	0.5 mL	No	25 g 1 inch

🌒 Vaccines with similar names that may lead to vaccine errors. 🛛 🚺 Products are subject to change based on availability.



### Catch-up (Resuming or Starting Immunizations)

- Disruptions in vaccine schedules do not require that a series be re-started, regardless of the length of time between doses.
- For catch-up schedules, please refer to the Canadian Immunization Guide for age-specific recommendations:

# Additional Publicly Funded Vaccines for Patients at High-Risk of Vaccine-preventable Disease

Patients at high-risk of vaccine-preventable diseases are eligible for

other vaccines based on condition-specific guidance.

nova scotia health Public Health

Updated August 2023

- Less than 7 years old; 7 - 17 years old; 18 years or older



## **Routine Vaccine Administration Guide**

i Please refer to the DHW Publicly Funded Vaccine Schedule here

## **Publicly Funded Vaccines in Nova Scotia**

(Childhood, School, Adult) Starting in Infancy (By Age)

CHILDHOOD VACCINES	VACCINE	PRODUCT NAME	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZ	E		
2 MONTHS	DTaP-IPV-Hib 🌗	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g	2 separate needles at least 1-inch apart		
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	1-inch			
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicab	le		
	DTaP-IPV-Hib リ	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g	2 separate		
<b>4</b> молтнs	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	1-inch l	needles at least 1-inch apart		
	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicab	le		
6	DTaP-IPV-Hib リ	Pediacel	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g 1-inch			
монтня	Rotavirus	RotaTeq	Orally administered	Oral	2 mL	No	Not applicable			
	Pneumococcal conjugate (Pneu-C-13)	Prevnar 13	Vastus lateralis (thigh)	Intramuscular	0.5 mL	No	25 g	2 separate needles at least 1-inch apart		
<b>12</b> MONTHS	Meningococcal group C conjugate	Neisvac-C Menjugate	Vastus lateralis (thigh)	Intramuscular	0.5 mL	NeisVac-C (No) Menjugate (Yes)	1-inch			
	MMRV (measles, mumps, rubella, varicella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine- specific diluent)	25 g 5/8- inch			
18	DTaP-IPV-Hib 🌗	Pediacel	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch			
MONTHS	MMRV (measles, mumps, rubella, vari-cella)	Priorix Tetra ProQuad	Upper arm	Subcutaneous	0.5 mL	Yes (vaccine- specific diluent)	25 g 5/8-inch			
<b>4-6</b> years	Tdap-IPV 🌗	Adacel Polio Boostrix Polio	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch			
SCHOOL VACCINES - Catch-up vaccines available through Public Health and/or Primary Care Provider prior to 19th Birthday										
	HPV series (0, 6 months)	Gardasil 9	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch			
GRADE 7 SCHOOL VACCINES	Hepatitis B series (0, 6 months)	Engerix-B Recombivax	Deltoid	Intramuscular	1.0 mL	No	25 g 1-inch			
	Tdap	Adacel Boostrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch			
	Meningococcal Quadrivalent (ACYW135)	Menveo Menactra Nimenrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch			

U vaccines with similar names that may lead to vaccine errors. i Products are subject to change based on availability.

**Routine Vaccine Administration Guide i** Please refer to the DHW Publicly Funded Vaccine Schedule **here** 

**Publicly Funded Vaccines in Nova Scotia** 

(Childhood, School, Adult) Starting in Infancy (By Age)

ADULT VACCINES	VACCINE	PRODUCT NAME	SITE	ROUTE	DOSE	RECONSTITUTION	NEEDLE SIZE
EVERY <b>10</b> YEARS	Td (Booster)	Td absorbed	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
ONCE AS AN ADULT	Tdap *If not previously given in adulthood. May receive one dose 10 years after receiving a tetanus containing vaccine.	Adacel Boostrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
PREGNANT PERSONS	Tdap: offered every pregnancy between 27 and 32 weeks	Adacel Boostrix	Deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
ADULTS BORN IN 1970 OR LATER	MMR (Booster)	MMRII (SC only) Priorix (SC preferred)	SC: Upper Arm IM: Deltoid	Subcutaneous Intramuscular	0.5 mL	Yes (vaccine- specific diluent)	SC: 25g 5/8-inch IM: 25g 1-inch
VACCINES FOR SENIORS							
65 YEARS AND OLDER	Pneumococcal polysaccharide (Pneu-P-23)	Pneumovax 23	IM: Deltoid SC: Upper Arm	Intramuscular Subcutaneous	0.5 mL	No	IM: 25 g 1-inch SC: 25 g 5/8-inch
65 YEARS AND OLDER	High-dose influenza; annually	Fluzone HD	Deltoid	Intramuscular	0.7 mL	No	25 g 1-inch
VACCINES FOR ALL AGES							
ANNUALLY, FROM 6 MONTHS OF AGE ONWARD	Influenza 1st-time recipients aged 6 months – 9 years: 2 doses (0, 4 weeks)	Fluzone Quad Flulaval Tetra Afluria Tetra (>5 years old only)	6-12 months old: vastus lateralis (thigh) Older than 12 months: deltoid	Intramuscular	0.5 mL	No	25 g 1-inch
i) For information on vaccination and pregnancy see here. i) For information on Covid-19 vaccination see here. i) Products are subject to change based on availability.							

#### Catch-up (Resuming or Starting Immunizations)

- Disruptions in vaccine schedules do not require that a series be re-started, regardless of the length of time between doses.
- For catch-up schedules, please refer to the Canadian Immunization Guide for age-specific recommendations:

## Additional Publicly Funded Vaccines for Patients at High-Risk of Vaccine-preventable Disease

Patients at high-risk of vaccine-preventable diseases are eligible for other vaccines based on **condition-specific guidance.** 



- Less than 7 years old; 7 - 17 years old; 18 years or older